

Cat. No. DSIU002

Vero (kidney, African green monkey, *Cercopithecus aethiops*)

Culture Medium :

MEM · E + 2mM Glutamine + 10% FBS

Cell line description :

The Vero cell line was initiated from the kidney of a normal adult African green monkey on March 27, 1962, by Y. Yasumura and Y. Kawakita at the Chiba University in Chiba, Japan (Nippon Rinsho 21: 1209, 1963).

The Vero cell line has been employed extensively in virus replication studies and plaque assays. The cell line has been useful for assay of SV-40, SV-5, measles, arboviruses, reoviruses, rubella, simian adenoviruses and polioviruses (Kitasato Arch. Exp. Med. 37:27, 1964; Bact. Proc., P.112, 1965; Proc. Soc. Exp. Bio. Med. 125: 119, 1967; *ibid.*, 125: 602, 1967; *ibid.*, 125: 636, 1967; *ibid.*, 125: 741, 1967; *ibid.*, 127: 642, 1968; Nature(Lond.) 216: 271, 1967; Arch. Gesamte Virusforsch. 21: 155-169, 1967; *ibid.*, 21: 243-252, 1967).

Freeze Medium : TC-Protector(Cell Freezing Media)

Morphology : Fibroblast-like

Split Ratio : 1:2 – 1:6

Virus Susceptibility : Simian Adenovirus, Dengue Virus, Herpes Simplex Virus, Varicella Zoster Cytomegalovirus, Measles Virus, Mumps Virus, Respiratory Syncytial Virus, Newcastle Disease Virus, Canine Distemper Virus, Simian Virus Type5, Subacute Sclerosing Panencephalitis Virus, Simian Virus Type40, Poliovirus Type3, Coxsackie TypeB, Echovirus, Reovirus Type1, Reovirus Type2, Reovirus Type3, Rubella Virus

株式会社 ケー・エー・シー

試薬事業部

〒661-0978 兵庫県尼崎市久々知西町2丁目1-20

(お問合せ先)

TEL : 06-6435-9747 FAX : 06-6435-9748

URL : <https://www.kacnet.co.jp/>

E-mail : shiyaku-info@kacnet.co.jp